# Package: HGDMr (via r-universe)

February 22, 2025

Type Package

Title Hysteretic and Gatekeeping Depressions Model

Version 1.0.0

Date 2025-02-19

Author Kevin Shook [cre, aut]

Maintainer Kevin Shook <kevin.shook@usask.ca>

Description Implementation of the Hysteretic and Gatekeeping Depressions Model (HGDM) which calculates variable connected/contributing areas and resulting discharge volumes in prairie basins dominated by depressions (``slough" or ``potholes"). The small depressions are combined into a single ``meta" depression which explicitly models the hysteresis between the storage of water and the connected/contributing areas of the depressions. The largest (greater than 5% of the total depressional area) depression (if it exists) is represented separately to model its gatekeeping, i.e. the blocking of upstream flows until it is filled. The methodolgy is described in detail in Shook and Pomeroy (2025, <doi:10.1016/j.jhydrol.2025.132821>).

License GPL-3

URL https://github.com/CentreForHydrology/HGDMr

**Depends** R (>= 4.0.0)

Imports stringr, stats

Suggests knitr, testthat, rmarkdown, readr, ggplot2

VignetteBuilder knitr

LazyData true

**Encoding** UTF-8

RoxygenNote 7.3.2

Config/pak/sysreqs libicu-dev

Repository https://centreforhydrology.r-universe.dev

2

RemoteUrl https://github.com/centreforhydrology/hgdmr RemoteRef HEAD RemoteSha db262b630604138a69b0b75734d9764cba52c41a

# Contents

HGDMr-package			2
A	•	4	
			7

# Index

HGDMr-package

The Hysteretic and Gatekeeping Depressions Model (HGDM)

# Description

**HGDMr** is an implementation of the Hysteretic and Gatekeeping Depressions Model (HGDM) which is a one-dimensional model of the effects of varying connected/contributing fractions of Canadian Prairie basins.

Using specified fluxes (rainfall, snowmelt, upland runoff and evaporation), HGDM computes a) the time-varying connected/contributing fraction of a basin having depressional storage and b) the depth of discharge in each time interval.

Note that this model does not route the flows through the basin. This would require the use of another R package, such as **RHMS**, which has the function 'reachRouting'.

#### Author(s)

Maintainer: Kevin Shook <kevin.shook@usask.ca>

#### References

Shook, Kevin R., and John W. Pomeroy. "The Hysteretic and Gatekeeping Depressions Model - A New Model for Variable Connected Fractions of Prairie Basins." Journal of Hydrology 654 (June 1, 2025): 132821. https://doi.org/10.1016/j.jhydrol.2025.132821.

Shook, Kevin R., Zhihua He, John W. Pomeroy, Chris Spence, and Colin J. Whitfield. "A Practitioner-Oriented Regional Hydrology Data Product for Use in Site-Specific Hydraulic Applications." Scientific Data 11, no. 1 (October 14, 2024): 1125. https://doi.org/10.1038/s41597-024-03962-1.

Clark, Martyn P., and Kevin R. Shook. "The Numerical Formulation of Simple Hysteretic Models to Simulate the Large-Scale Hydrological Impacts of Prairie Depressions." Water Resources Research 58, no. 12 (2022): e2022WR032694. https://doi.org/10.1029/2022WR032694.

Shook, Kevin, Simon Papalexiou, and John W. Pomeroy. "Quantifying the Effects of Prairie Depressional Storage Complexes on Drainage Basin Connectivity." Journal of Hydrology 593 (February 1, 2021): 125846. https://doi.org/10.1016/j.jhydrol.2020.125846.

#### daily\_7120951600

Shook, Kevin, John W Pomeroy, Christopher Spence, and Lyle Boychuk. "Storage Dynamics Simulations in Prairie Wetland Hydrology Models: Evaluation and Parameterization." Hydrological Processes 27, no. 13 (June 2013): 1875–89. https://doi.org/10.1002/hyp.9867.

#### See Also

Useful links:

https://github.com/CentreForHydrology/HGDMr

daily\_7120951600 daily basin 7120951600 PHyDAP fluxes

#### Description

A dataframe of daily CRHM fluxes modelled for basin 7120951600. The fluxes were taken from the PHyDAP project https://www.frdr-dfdr.ca/repo/dataset/7ce4bd7a-4bcc-4f8c-8129-32a691f46c8e hourly outputs of CRHM models forced with ERA5 data over the period 1950-2020. The fluxes were then aggregated to daily values.

#### Usage

daily\_7120951600

#### Format

A dateframe with 25932 rows and 5 columns spanning the period 1950-2020.

#### Details

Variables:

date R date

**rainfall** Daily rainfall on water (mm)

snowmelt Daily snow melt on water (mm)

runoff Daily upland runoff (mm)

evap Daily water evaporation (mm)

#### Source

PHyDAP

#### References

Shook, Kevin R., Zhihua He, John W. Pomeroy, Chris Spence, and Colin J. Whitfield. "A Practitioner-Oriented Regional Hydrology Data Product for Use in Site-Specific Hydraulic Applications." Scientific Data 11, no. 1 (October 14, 2024): 1125. https://doi.org/10.1038/s41597-024-03962-1.

## Description

Applies the Hysteretic and Gatekeeping Depressions Model to basin-scale fluxes determined by hydrological modelling to calculate the outflows during a given time interval. Note than no routing is performed.

## Usage

```
HGDM(
  upland_area = NULL,
  small_depression_area = NULL,
  large_depression_area = NULL,
  area_units = "km2",
 max_small_depression_storage = 0,
 max_large_depression_storage = 0,
  initial_small_depression_storage = 0,
  initial_large_depression_storage = 0,
  storage_units = "mm",
  small_depressions_initial_connected_fraction = 0,
  upland_fraction_to_small = 0,
  upland_fraction_to_large = 0,
  upland_fraction_to_outlet = 0,
  small_fraction_to_large = 0,
  forcings = NULL,
  small_p = NULL,
  large_rating = 0,
  sub_intervals = 1
```

#### Arguments

)

upland_area	Required. Area of uplands, which drain to the outlet, small depressions or the large depression.			
small_depression	on_area			
	Required. Area of small depressions.			
large_depression	on_area			
	Optional. If 0 or NULL large depression is not \ modelled.			
area_units	Units of all areas. Must be one of 'km2' (default), 'ha' or 'm2'.			
<pre>max_small_depression_storage</pre>				
	Maximum depth of storage in small depressions.			
<pre>max_large_depre</pre>	ession_storage			
	Maximum depth of storage in large depressions.			

#### HGDM

initial_small_d	epression_storage			
	Initial depth of storage in small depressions.			
initial_large_depression_storage				
	Initial depth of storage in large depressions.			
storage_units	Units of all storage depths. Must be one of 'mm' (default) 'm', or 'm3'. If a depth is specified then it will be converted to a volume by multiplying by the appropriate area.			
<pre>small_depressions_initial_connected_fraction</pre>				
	Initial connected fraction (0-1).			
upland_fraction_to_small				
	Fraction of uplands draining to small depressions. If 0 then the small depressions are unlikely to fill.			
upland_fraction_to_large				
	Fraction of uplands draining to large depression. This is the basin of the large depression.			
upland_fraction_to_outlet				
	Fraction of uplands draining directly to outlet. Analogous to the effective fraction.			
<pre>small_fraction_to_large</pre>				
	Fraction of small depression area draining into large depression. Governed by location of large depression in the basin.			
forcings	Required. A data frame of time series of rainfall, snowmelt, evap, and runoff. The first variable must be either date (an ${\bf R}$ date) or datetime (a POSIXct date-time).			
small_p	Parameter for small depression water volume-area relationship.			
large_rating	Rating curve parameters for large depression.			
<pre>sub_intervals</pre>	Number of sub-intervals for solution of each time step.			

# Value

Returns a data frame. Depending on whether or not a large depression was specified, the data frame will have differing variables. Note that regardless of the units specified for areas and volumes, all of the variables returned are in SI dimensions, i.e. 'm' and ' $m^3/s'$  values

If no large depression is specified, the returned variables are:

date or datetime R date or POSIXct datetime.

**total\_contrib\_frac** The connected/contributing fraction of the basin. Includes both the meta depression and the upland fraction connected to the outlet.

total\_outflow\_volume The volume of outflow (m<sup>3</sup>) in the interval.

small\_depression\_contrib\_frac The connected/contributing fraction of the meta depression.

small\_depression\_water\_volume The volume of water (m<sup>3</sup>) retained in the meta depression.

small\_depression\_water\_depth The depth of water (m) retained in the meta depression.

small\_depression\_water\_area The area of water (m<sup>2</sup>) retained in the meta depression.

If there is a large depression, then 'total\_contrib\_frac' includes the effect of the large depression and the additional variables are also returned:

date or datetime R date or POSIXct datetime.

**large\_depression\_contrib\_frac** The connected/contributing fraction of the large depression. **large\_depression\_water\_volume** The volume of water (m<sup>3</sup>) retained in the large depression. **large\_depression\_water\_area** The area of water (m<sup>2</sup>) retained in the large depression.

#### Examples

```
{
daily_fluxes <- daily_7120951600</pre>
basin_area <- 100</pre>
small_depression_frac <- 0.24</pre>
small_depression_area <- small_depression_frac * basin_area</pre>
large_depression_area <- 0</pre>
upland_area <- basin_area - (small_depression_area + large_depression_area)</pre>
area_units <- "km2"
max_small_depression_storage <- 300</pre>
max_large_depression_storage <- 0</pre>
initial_small_depression_storage <- max_small_depression_storage / 2</pre>
initial_large_depression_storage <- max_large_depression_storage / 2</pre>
storage_units <- "mm"</pre>
small_depressions_initial_connected_fraction <- 0</pre>
upland_fraction_to_small <- 0.98
upland_fraction_to_large <- 0
upland_fraction_to_outlet <- 0.02</pre>
small_fraction_to_large <- 0</pre>
small_p <- 1.2
large_rating <- 1.4</pre>
sub_intervals <- 1</pre>
results <- HGDM(upland_area,
small_depression_area,
large_depression_area = 0,
area_units = "km2", max_small_depression_storage,
max_large_depression_storage,
initial_small_depression_storage,
initial_large_depression_storage,
storage_units,
small_depressions_initial_connected_fraction,
upland_fraction_to_small,
upland_fraction_to_large,
upland_fraction_to_outlet,
small_fraction_to_large,
forcings = daily_fluxes[1:100,],
small_p = small_p,
large_rating = large_rating,
sub_intervals = sub_intervals)
}
```

6

# Index

HGDMr-package, 2

daily\_7120951600,3

HGDM, 4 HGDMr (HGDMr-package), 2 HGDMr-package, 2